



PHOTO CREDIT: Dr. William M. Johnson

Many Gulf Coast gardeners think of gardening as a spring and summer activity. Eating carrots right out of the winter garden is a real treat. Once they are kissed by a frost, they begin to sweeten up.

Many Gulf Coast gardeners think of gardening as a spring and summer activity. As I walked through the Master Gardener Discovery Garden in Carbine Park last week on a very cool and misty Thursday morning, I was inspired by the abundance of winter vegetables being grown — broccoli, cauliflower, kale, beets, Irish potatoes, cab-

bage, onions, and the list goes on.

Many of the vegetables we planted in late summer and early fall are now ready to harvest — or will be soon. It is important to harvest vegetables at the proper stage for best results, so here are a few guidelines for some common cool-

season crops.

Root crops are usually harvested when the top of the root becomes readily visible at ground-level but it is easy enough to brush aside the soil at the base of the leaves to check on the size of the root. Harvest radishes and carrots when the root is about 1



December 13, 2017



Dr. William Johnson is a horticulturist with the Galveston County Office of Texas A&M AgriLife Extension Service. Visit his website at <http://aggie-horticulture.tamu.edu/galveston>.

News column printed in the Galveston Daily News, The Post, and other Galveston County Newspapers.

inch across.

Carrots can be left in the ground once they are mature and can be harvested as needed, and the tops can be used as a parsley substitute. Turnips should be harvested when they are 2 to 3 inches in diameter, and rutabagas (a close relative) when they are 4 or 5 inches in diameter. Beets are best harvested at 2 to 3 inches and parsnips at 1 1/2 to 2 inches.

Broccoli heads are not harvested based on the size of the head, but when the largest individual flower buds are about the size of a kitchen match head. Cut the primary crown (where the individual heads come together) when it's about 4 inches across. Do not allow the heads to remain on the plant so long that some of the buds open to produce a yellow flower. Remember that smaller side heads will develop after the main head has been harvested, so leave the plant in place for additional harvest.

Harvesting cauliflower also depends more on the appearance of the head rather than its size. The curds of the head should be relatively smooth, very much like the cauliflower that you buy in the supermarket. If allowed to stay on the plant too long, the head will begin to separate and lose quality. If you did not blanch your cauliflower by covering the head with the plant's leaves, it may have a purple, green or yellow tint to it. This does not greatly affect the quality of the head.

harvested frequently by breaking off the lowest, largest leaves (this is called cropping). Harvest the entire head of semi-heading varieties of lettuce such as bibb, buttercrunch and romaine when the head is fully developed.

Cabbage is ready to harvest when the head is solid and hard. Cabbage is one of the few crops that may be left in the garden after they are ready to harvest, although the heads may split. If you are going to leave fully formed heads in the garden, rotate the entire plant one-half turn to prevent splitting (this slows water uptake by breaking some of the roots).

Snow peas and edible podded peas are productive, delicious and well worth growing. Harvest snow peas when the pods are full size but still quite flat. Edible-podded peas, such as sugar snap peas, should be harvested when the pods are full and round but before the peas inside the pod have fully developed. Both types of peas should be checked daily and harvested frequently.

Cold protection for winter vegetables

Last week started out with daytime temperatures in the low eighties on Monday and ended up on Thursday and Friday with snow and freezing weather. Although winter vegetables are generally hardy, new plantings may need to be protected from

Harvesting Tips for Winter Vegetables

hard freezes as will certain vegetables near or at harvest stage.

If temperatures below 30 degrees are predicted, young seedlings should receive special attention by completely covering them with a 4-to-6-inch layer of loose mulch like leaves or pine straw. The mulch may remain over the plants for a few days, but remove it as soon as the freezing episode is over. Other protections include plastic coverings supported to keep them off the plants, fabric sheets or floating row coverings may also be used.

The following lists will give you a quick guide to the ability of some vegetables to endure freezes.

- Protect or harvest the following vegetables if temperatures are predicted to go below 30 degrees: broccoli, cauliflower, lettuce and peas.

- Vegetables that can tolerate temperatures down to the mid-20s with little or no damage: Swiss chard, Chinese cabbage, kohlrabi, mustard, spinach, radishes and turnips (although the leaves of radishes and turnips are moderately hardy, their roots are very hardy).

- Vegetables that can tolerate temperatures in the low 20s and even the teens, especially if given some protection: beets, Brussels sprouts, carrots, celery, collards, garlic, onions, parsley, leeks and shallots.